

# New explicit examples of conformally constrained Willmore minimizers

Reiner Schätzle<sup>1</sup>

Fachbereich Mathematik der Eberhard-Karls-Universität Tübingen,  
Auf der Morgenstelle 10, D-72076 Tübingen, Germany,  
email: schaeetz@everest.mathematik.uni-tuebingen.de

**Abstract:** By estimates of Li, Yau and Montiel, Ros, the Clifford torus  $T_{Cliff} := (1/\sqrt{2})(S^1 \times S^1) \subseteq S^3$  minimizes the Willmore energy in its conformal class. We extend this to the flat constant mean curvature tori  $T_r := rS^1 \times \sqrt{1-r^2}S^1 \subseteq S^3$  for  $r \approx 1/\sqrt{2}$ , thereby getting new explicit examples of conformally constrained Willmore minimizers. Actually the minimization of  $T_r$  is slightly stronger, and we obtain supplementing the estimates of Li, Yau and Montiel, Ros, that the Clifford torus minimizes the Willmore energy in an open neighbourhood of its conformal class, where the neighbourhood may depend on the codimension.

**Keywords:** Willmore surfaces, conformal parametrization.

**AMS Subject Classification:** 53 A 05, 53 A 30, 53 C 21.

---

<sup>1</sup>This work was supported by the DFG Sonderforschungsbereich TR 71 Freiburg - Tübingen. It is joint work with Cheikh Birahim Ndiaye, Tübingen.