УДК 539.12

ON HIGHER REGULARITY AND MONOIDAL CATEGORIES

S. A. Duplij[1)], **W. Marcinek**[2)]

[1)] *Department of Physics and Technology, V. N. Karazin Kharkov National University, Kharkov 61077, Ukraine*
E-mail: Steven.A.Duplij@univer.kharkov.ua. Internet: http://gluon.physik.uni-kl.de/~duplij

[2)] *Institute of Theoretical Physics, University of Wrocław, Pl. Maxa Borna 9, 50-204 Wrocław, Poland*

Received May 12, 2000

In general abstract algebraic language we extend "invertibility" to "regularity" for categories. Higher regularity conditions and "semicommutative" diagrams are introduced. Distinction between commutative and "semicommutative" cases is measured by non-zero obstruction proportional to the difference of some self-mappings (obstructors) $e^{(n)}$ from the identity, which allows us to generalize the notion of functor and to "regularize" braidings and related structures in monoidal categories. Also we propose a "noninvertible" analog of the Yang-Baxter equation.
KEYWORDS : morphism, regularity, obstruction, monoidal category, functor, Yang-Baxter equation

The concept of regularity was introduced by von Neumann [1] and applied by Penrose to matrices [2]. After that study of regularity was developed in many different fields, e.g. generalized inverses theory [3] and semigroup theory [4]. We consider here this concept in categorical language [5] and introduce the most abstract form of higher regularity conditions (firstly introduced in [6]).

Let X and Y be two arbitrary sets. A mapping f from X to Y is defined by a prescription which assigns an element of Y to each element of X, i.e. $f : X \to Y$. Injective mapping (injection) assigns different images to different elements, and in surjective mapping (surjection) every image has at least one pre-image. Bijection has both properties. Usually inverse mapping f^{-1} is defined as a new mapping $g : Y \to X$ which assigns to each $y \in Y$ such $x \in X$ that $f(x) = y$ and so $f^{-1} = g$. For injective f and any $A \subset X$ it is imposed the following "invertibility" condition

$$f^{-1}(f(A)) = A. \tag{1}$$

For surjective f and $B \subset Y$ the standard "invertibility" condition is

$$f\left(f^{-1}(B)\right) = B. \tag{2}$$

These conditions are strong, because they imply possibility to solve the equation $f(x) = y$ for all elements. In many cases, especially while considering supersymmetric theories, there naturally appear noninvertible morphisms [6, 7] and semigroups [8, 9]. That obviously needs extending some general assumptions. We propose to extend the "invertibility" conditions (1)–(2) in the following way (which comes from analogy of regularity in semigroup theory [4]). We introduce less restricted "regular" f^* mapping by extending "invertibility" to "regularity" in following way

$$f(f^*(f(A))) = f(A). \tag{3}$$

For the second equation (2) we have the "reflexive regularity" condition

$$f^*(f(f^*(B))) = f^*(B). \tag{4}$$

REGULAR MORPHISMS

We distinguish among all mappings $X \to Y$ the morphisms satisfying closure and associativity. That defines a category $\mathfrak{C}$ with objects $\mathrm{Ob}\,\mathfrak{C}$ as sets X, Y, Z and morphisms $\mathrm{Mor}\,\mathfrak{C}$ as mappings $f : X \to Y$ between them (or $f = \mathrm{Mor}(X, Y)$) [5]. For composition $h : X \to Z$ of morphisms $f : X \to Y$ and $g : Y \to Z$ instead of $h(x) = g(f(x))$ for mappings we use the notation $h = g \circ f$. Associativity implies that $h \circ (g \circ f) = (h \circ g) \circ f = h \circ g \circ f$. Let us consider "invertibility" properties of morphisms in general. If f satisfies the "right invertibility" condition $f \circ f^{-1} = Id_Y$ for some $f^{-1} : Y \to X$ then f is called a *retraction*, and if f satisfies the "left invertibility" condition $f^{-1} \circ f = Id_X$, then it is called a *coretraction*, where Id_X and Id_Y are identity mappings $Id_X : X \to X$ and $Id_Y : Y \to Y$ for which $\forall x \in X$, $Id_X(x) = x$ and $\forall y \in Y$, $Id_Y(y) = y$. These requirements sometimes are very strong to be fulfilled (see e.g. [10]). To obtain more weak conditions one has to introduced the following "regularity" conditions

$$f \circ f^*_{in} \circ f = f, \tag{5}$$

where f^*_{in} is called an *inner inverse* [3], and such f is called *regular*. Similar "reflexive regularity" conditions

$$f^*_{out} \circ f \circ f^*_{out} = f^*_{out} \tag{6}$$

defines an *outer inverse* f^*_{out}. Notice that in general $f^*_{in} \neq f^*_{out} \neq f^{-1}$ or it can be that f^{-1} does not exist at all. If f^*_{in} is an inner inverse, then

$$f^* = f^*_{in} \circ f \circ f^*_{in} \tag{7}$$

is always both inner and outer inverse or *generalized inverse* (quasi-inverse) [3], and so for any regular f there exists (need not be unique) f^* from (7) for which both regularity conditions (5) and (6) hold. Let us consider a composition of two morphisms and its "invertibility" properties. It can be shown, that a retraction is an epimorphism, and a regular monomorphism is a coretraction [3]. If the composition $h = g \circ f$ belongs to the same class of functions (closure), then all such morphisms form a semigroup of such functions [4]. If for any $f : X \to Y$ there will be a unique $f^* : Y \to X$ satisfying (5)–(6), this semigroup is called an inverse semigroup [4] which we denote $\mathfrak{F}$.

Let us define two idempotent "projection operators" $\mathcal{P}_f = f \circ f^*$, $\mathcal{P}_f : Y \to Y$ and $\mathcal{P}_{f^*} = f^* \circ f$, $\mathcal{P}_{f^*} : X \to X$ satisfying $\mathcal{P}_f \circ \mathcal{P}_f = \mathcal{P}_f$, $\mathcal{P}_f \circ f = f \circ \mathcal{P}_{f^*} = f$ and $\mathcal{P}_{f^*} \circ \mathcal{P}_{f^*} = \mathcal{P}_{f^*}$, $\mathcal{P}_{f^*} \circ f^* = f^* \circ \mathcal{P}_f = f^*$. If we introduce the $*$-operation $(f)^* = f^*$ by formulas (5)–(6) and assume that this operation acts on the product of morphisms $f : X \to Y$ and $g : Y \to Z$ in the following way $(g \circ f)^* = f^* \circ g^*$, then commutativity of projectors $\mathcal{P}_f \circ \mathcal{P}_{g^*} = \mathcal{P}_{g^*} \circ \mathcal{P}_f$ leads to closure of the semigroup, i.e. the product $g \circ f$ also satisfies both regularity conditions (5)–(6).

HIGHER REGULARITY

Let us introduce higher analogs of regularity conditions (5)–(6) (they were proposed for some particular case (noninvertible analogs of supermanifolds) in [6,7]). Let we have two elements f and its regular f^* (in sense of (5)) of the semigroup $\mathfrak{F}$. Consider a third morphism $f^{**} : X \to Y$ and analyze the action $f \circ f^* \circ f^{**} : X \to Y$. This means the composition $f \circ f^* \circ f^{**}$ cannot be equal to identity Id_X. Therefore it is possible to "regularize" $f \circ f^* \circ f^{**}$ in the following way

$$f \circ f^* \circ f^{**} \circ f^* = f^*. \tag{8}$$

This formula can be called as 2-regularity condition and be considered as a definition of $**$-operation. For 3-regularity and $f^{***} : Y \to X$ we can obtain an analog of (5) in the form

$$f \circ f^* \circ f^{**} \circ f^{***} \circ f = f. \tag{9}$$

By recursive considerations we can propose the following formula of n-regularity

$$f \circ f^* \circ f^{**} \ldots \circ \overbrace{f^{* * \cdots *}}^{2k} \circ f^* = f^*, \quad f \circ f^* \circ f^{**} \ldots \circ \overbrace{f^{* * \cdots *}}^{2k+1} \circ f = f. \tag{10}$$

Note that for even number of stars $\overbrace{f^{* * \cdots *}}^{2k} : X \to Y$ and for odd number of stars $\overbrace{f^{* * \cdots *}}^{2k+1} : Y \to X$. We introduce "higher projector" by the formula

$$\mathcal{P}^{(n)}_f = f \circ f^* \circ f^{**} \ldots \circ \overbrace{f^{* * \cdots *}}^{n}. \tag{11}$$

It is easy to check the following properties

$$\mathcal{P}^{(2k)}_f \circ f^* = f^*, \quad \mathcal{P}^{(2k+1)}_f \circ f = f. \tag{12}$$

and idempotence $\mathcal{P}^{(n)}_f \circ \mathcal{P}^{(n)}_f = \mathcal{P}^{(n)}_f$.

SEMICOMMUTATIVE DIAGRAMS AND OBSTRUCTORS

Obviously, that for two morphisms $f : X \to Y$ and $g : Y \to X$ instead of "invertibility" $g \circ f = Id_X$ we have the same generalization as regularity (5), i.e. $f \circ g \circ f = f$, where g plays the role of an inner inverse [3].



Invertible morphisms Noninvertible (regular) morphisms

Usually, for 3 objects X, Y, Z and 3 morphisms $f : X \to Y$ and $g : Y \to Z$ and $h : Z \to X$ one can have the "invertible" triangle commutative diagram $h \circ g \circ f = Id_X$. Its regular extension has the form

$$f \circ h \circ g \circ f = f. \tag{13}$$

Such a diagram (from the right)



can be called a *semicommutative diagram*. This triangle case can be expanded on any number of objects and morphisms. To measure difference between semicommutative and commutative cases let us introduce self-mappings $e_X^{(n)} : X \to X$ which are defined by

$$e_X^{(1)} = Id_X,\ e_X^{(2)} = g \circ f,\ e_X^{(3)} = h \circ g \circ f,\ \ \ldots \tag{14}$$

It is obvious that for commutative diagrams all $e_X^{(n)}$ are equal to identity $e_X^{(n)} = Id_X$. The deviation of $e_X^{(n)}$ from identity will give us measure of obstruction of commutativity, and therefore we call $e_X^{(n)}$ *obstructors*. The minimum number $n = n_{obstr}$ for which $e_X^{(n)} \neq Id_X$ occurs will define a quantitative measure of obstruction n_{obstr}. In terms of obstructors $e_X^{(n)}$ the n-regularity condition can be written in the short form $f \circ e_X^{(n)} = f$. From this equation and definitions (14) it simply follows that obstructors $e_X^{(n)}$ are idempotents.

REGULARIZATION OF MONOIDAL CATEGORIES

Let $\mathfrak{C}$ be a monoidal category equipped with a monoidal operation $\otimes : \mathfrak{C} \times \mathfrak{C} \longrightarrow \mathfrak{C}$. A triple of objects X, Y, Z is said to be a regular 3-cycle if and only if every sequence of morphisms $X \stackrel{f}{\longrightarrow} Y \stackrel{g}{\longrightarrow} Z \stackrel{h}{\longrightarrow} X$ define uniquely the morphism $e_X^{(3)} : X \longrightarrow X$ by the following relation $e_X^{(3)} := h \circ g \circ f$ and subjects to the relation $f \circ h \circ g \circ f = f$. The object Y is said to be a *(first) regular dual* of X, and the object Z is called the *second regular dual* of X. We denote by $C_3(\mathfrak{C})$ the collection of all regular 3-cycles on $\mathfrak{C}$. This collection is said to be *regularity* in $\mathfrak{C}$. The generalization to arbitrary $n \geq 4$ is obvious. Let X, Y, Z and X', Y', Z' be two regular 3-cycles in $\mathfrak{C}$. Then the morphism $f : X \longrightarrow X'$ such that $f \circ e_X^{(3)} = e_{X'}^{(3)} \circ f$ is said to be a *3-cycle morphism*. If $f : X \longrightarrow X'$ and $g : X' \longrightarrow X''$ are two 3-cycle morphisms, then the composition $g \circ f : X \to X''$ is also a 3-cycle morphism. Moreover the regularity $C_3(\mathfrak{C})$ forms a monoidal category with 3-cycles as objects and 3-cycle morphisms. The monoidal product of two regular 3-triples X, Y, Z and X', Y', Z' is the triple $X \otimes X', Y \otimes Y', Z \otimes Z'$ which is also a regular 3-cycle. The category $C_3(\mathfrak{C})$ is said to be a *regularization* of $\mathfrak{C}$.

The morphisms $e_X^{(n)}$ can be used to extend the notion of a functor $\mathsf{F} : \mathfrak{C}_1 \to C_2$. All the standard definitions of a functor (as a mapping of one category to another with preserving composition of morphisms [5]) do not changed, but preservation of identity $\mathsf{F}(Id_{X_1}) = Id_{X_2}$, where $X_2 = \mathsf{F}X_1$, $X_1 \in \mathrm{Ob}\mathfrak{C}_1$, $X_2 \in \mathrm{Ob}\mathfrak{C}_2$, can be replaced by requirement of preservation of morphisms $e_X^{(n)}$ as

$$\mathsf{F}^{(n)}\left(e_{X_1}^{(n)}\right) = e_{X_2}^{(n)}, \tag{15}$$

where $e_{X_1}^{(n)} \in \mathrm{Mor}\mathfrak{C}_1$, $e_{X_2}^{(n)} \in \mathrm{Mor}\mathfrak{C}_2$ defined in (14) for two categories. Then the generalized functor $\mathsf{F}^{(n)}$ becomes n-dependent. From (14) it follows that $n = 1$ corresponds to the standard functor, i.e. $\mathsf{F}^{(1)} = \mathsf{F}$.

HIGHER REGULAR YANG-BAXTER EQUATION

Let us consider a symmetric monoidal category $\mathfrak{C}$ [5] playing an important role in quantum groups [11] and quantum statistics [12].In $\mathfrak{C}$ for any two objects X and Y and the operation $X \otimes Y$ one usually defines a natural isomorphism ("braiding" [13]) by $\mathrm{B}_{X,Y} : X \otimes Y \to Y \otimes X$ satisfying the symmetry condition ("invertibility")

$$\mathrm{B}_{Y,X} \circ \mathrm{B}_{X,Y} = Id_{X \otimes Y} \tag{16}$$

which formally defines $\mathrm{B}_{Y,X} = \mathrm{B}_{X,Y}^{-1} : Y \otimes X \to X \otimes Y$. Note that possible nonsymmetric braiding in context of the noncommutative geometry was considered in [14]. Here we introduce a "regular" extension of the symmetry condition (16) in the form

$$\mathrm{B}_{X,Y} \circ \mathrm{B}_{X,Y}^{*} \circ \mathrm{B}_{X,Y} = \mathrm{B}_{X,Y}, \tag{17}$$

where in general $\mathrm{B}_{X,Y}^{*} \neq \mathrm{B}_{X,Y}^{-1}$. Such a category can be called a "regular" category to distinct from symmetric and "braided" categories [13].

In categorical sense the prebraiding relations usually are defined as [13, 14]

$$\mathrm{B}_{X\otimes Y,Z} = \mathbf{B}^R_{X,Z,Y}\circ\mathbf{B}^L_{X,Y,Z},\quad \mathrm{B}_{Z,X\otimes Y} = \mathbf{B}^L_{X,Z,Y}\circ\mathbf{B}^R_{X,Y,Z}, \tag{18}$$

$$\mathbf{B}^L_{X,Y,Z} = Id_X\otimes \mathrm{B}_{Y,Z},\quad \mathbf{B}^R_{X,Y,Z} = \mathrm{B}_{X,Y}\otimes Id_Z, \tag{19}$$

and prebraidings $\mathrm{B}_{X\otimes Y,Z}$ and $\mathrm{B}_{Z,X\otimes Y}$ satisfy (for symmetric case) the "invertibility" property $\mathrm{B}^{-1}_{X\otimes Y,Z}\circ \mathrm{B}_{X\otimes Y,Z} = Id_{X\otimes Y\otimes Z}$, where $\mathrm{B}^{-1}_{X\otimes Y,Z} = \mathrm{B}_{Z,X\otimes Y}$. In this notations the standard "invertible" Yang-Baxter equation is [11]

$$\mathbf{B}^R_{Y,Z,X}\circ\mathbf{B}^L_{Y,X,Z}\circ\mathbf{B}^R_{X,Y,Z} = \mathbf{B}^L_{Z,X,Y}\circ\mathbf{B}^R_{X,Z,Y}\circ\mathbf{B}^L_{X,Y,Z}. \tag{20}$$

Possible "noninvertible" (endomorphism semigroup) solutions of this equation without introduction of $e^{(n)}_X$ were studied in [15]. For "noninvertible" braidings satisfying regularity (17) it is naturally to exploit the obstructors $e^{(n)}_X$ instead of identity Id_X as

$$\mathbf{B}^{L(n)}_{X,Y,Z} = e^{(n)}_X\otimes \mathrm{B}_{Y,Z},\quad \mathbf{B}^{R(n)}_{X,Y,Z} = \mathrm{B}_{X,Y}\otimes e^{(n)}_Z, \tag{21}$$

to weaken prebraiding construction in the following way

$$\mathrm{B}^{(n)}_{X\otimes Y,Z} = \mathbf{B}^{R(n)}_{X,Z,Y}\circ\mathbf{B}^{L(n)}_{X,Y,Z},\quad \mathrm{B}^{(n)}_{Z,X\otimes Y} = \mathbf{B}^{L(n)}_{X,Z,Y}\circ\mathbf{B}^{R(n)}_{X,Y,Z}, \tag{22}$$

Then their "invertibility" can be also "regularized" as follows

$$\mathrm{B}^{(n)}_{X\otimes Y,Z}\circ\mathrm{B}^{(n)*}_{X\otimes Y,Z}\circ\mathrm{B}^{(n)}_{X\otimes Y,Z} = \mathrm{B}^{(n)}_{X\otimes Y,Z}, \tag{23}$$

where in general case $\mathrm{B}^{(n)*}_{X\otimes Y,Z}\neq \mathrm{B}^{-1}_{X\otimes Y,Z}$. Thus the corresponding n-"noninvertible" analog of the Yang-Baxter equation (20) is

$$\mathbf{B}^{R(n)}_{Y,Z,X}\circ\mathbf{B}^{L(n)}_{Y,X,Z}\circ\mathbf{B}^{R(n)}_{X,Y,Z} = \mathbf{B}^{L(n)}_{Z,X,Y}\circ\mathbf{B}^{R(n)}_{X,Z,Y}\circ\mathbf{B}^{L(n)}_{X,Y,Z}. \tag{24}$$

Its solutions can be found by application of the semigroup methods (see e.g. [15]). The introduced formalism can be used in analysis of categories with some weaken invertibility conditions, which can appear in nontrivial supersymmetric or noncommutative geometry constructions beyond the group theory.

Acknowledgments. One of the authors (S.D.) would like to thank Jerzy Lukierski for kind hospitality at the University of Wrocław, where this work was done, and to Andrzej Borowiec, Andrzej Frydryszak and Cezary Juszczak for valuable discussions and help during his stay in Wrocław.

REFERENCES

1. von Neumann J. // *On regular rings.* Proc. Nat. Acad. Sci. USA. 1936. V. **22**. P. 707-713.
2. Penrose R. // *A generalized inverse for matrices.* Math. Proc. Cambridge Phil. Soc. 1955. V. **51**. P. 406-413.
3. Nashed M. Z. Generalized Inverses and Applications. New York. *Academic Press*, 1976. 321 p.
4. Howie J. M. An Introduction to Semigroup Theory. London. *Academic Press*, 1976. 270 p.
5. MacLane S. Categories for the Working Mathematician. Berlin. *Springer-Verlag*, 1971. 189 p.
6. Duplij S. // *Noninvertibility and "semi-" analogs of (super) manifolds, fiber bundles and homotopies.* Kaiserslautern, 1996. 30 p. (*Preprint / Univ. Kaiserslautern*; KL-TH-96/10, q-alg/9609022).
7. Duplij S. // *On semi-supermanifolds.* Pure Math. Appl. 1998. V. **9**. № 3-4. P. 283-310.
8. Duplij S. // *On semigroup nature of superconformal symmetry.* J. Math. Phys. 1991. V. **32**. № 11. P. 2959-2965.
9. Duplij S. Semigroup methods in supersymmetric theories of elementary particles. Kharkov. *Habilitation Thesis, Kharkov State University*, math-ph/9910045, 1999. 483 p.
10. Davis D. L., Robinson D. W. // *Generalized inverses of morphisms.* Linear Algebra Appl. 1972. V. **5**. P. 329-338.
11. Majid S. // *Quasitriangular Hopf algebras and Yang-Baxter equations.* Int. J. Mod. Phys. 1990. V. **A5**. № 1. P. 1-91.
12. Marcinek W. // *Categories and quantum statistics.* Rep. Math. Phys. 1996. V. **38**. № 2. P. 149-174.
13. Joyal A., Street R. // *Braided monoidal categories.* North Ryde, New South Wales, 1986. 45 p. (*Preprint / Macquarie University*; Mathematics Reports 86008).
14. Marcinek W. // *Noncommutative geometry for arbitrary braidings.* J. Math. Phys. 1996. V. **35**. P. 2633-2641.
15. Li F. // *Weak Hopf algebras and new solutions of Yang-Baxter equation.* J. Algebra. 1998. V. **208**. № 1. P. 72-100.

О ВЫСШЕЙ РЕГУЛЯРНОСТИ И МОНОИДАЛЬНЫХ КАТЕГОРИЯХ

С. А. Дуплий[1], **В. Марчинек**[2]

[1] *Харьковский национальный университет им. В. Н. Каразина, пл. Свободы, 4, г. Харьков, 61077, Украина*
[2] *Институт теоретической физики, университет Вроцлава, пл. Макса Борна 9, 50-204 Вроцлав, Польша*

Мы расширяем "обратимость" на "регулярность" для категорий в абстрактном алгебраическом подходе. Введены условия высшей регулярности и "полукоммутативные" диаграммы. Различие между коммутативным и "полукоммутативным" случаями измеряется отличием некоторых отображений $e^{(n)}$ от единичного, что позволяет обобщить понятие функтора и "регуляризовать" подобные структуры в моноидальных категориях. Предложен также "необратимый" аналог уравнения Янга-Бакстера.

КЛЮЧЕВЫЕ СЛОВА: морфизм, регулярность, препятствие, моноидальная категория, функтор, уравнение Янга-Бакстера