

Montag, 13.05.2024 um 15:15 Uhr
R87, Wilhelm-Klemm-Str. 10

Polaromechanics: microcavity excitation-polaritons meet GHz phonons



© Dr. Alexander Kuznetsov

Dr. Alexander Kuznetsov

Paul-Drude-Institut für Festkörperelektronik, Berlin

Engineered coupling between optical, electronic and vibrational degrees of freedom opens a pathway to merge domains of cavity quantum electrodynamics and optomechanics. On the one hand, excitons can couple strongly to resonant photons, resulting in light-matter exciton-polariton quasiparticles. On the other hand, lattice phonons can couple efficiently to excitons via the deformation potential mechanism. Moreover, interactions between phonons, excitons and photons can be enhanced in semiconductor microcavities (MCs). Such hybrid systems allow to explore rich physics of coherent optomechanical interactions in non-linear driven-dissipative systems. In this talk, I will introduce the emerging field of polaromechanics [1], which studies interactions between non-equilibrium polariton Bose-Einstein-like condensates [2] and GHz phonons in hybrid phonon-photon (Al,Ga)As MCs [3]. I will describe, how we engineer MCs to enhance the coupling between polaritons and monochromatic GHz phonons that we can inject using piezoelectric transducers [4]. Then, I will discuss the adiabatic interaction between polaritons and phonons, i.e., when the polariton decoherence rate exceeds the phonon frequency, $\gamma_{\text{pol}} \gg \Omega_{\text{M}}$ [5]. I will then show that under strong optical pumping (large quasiparticle populations), polaritons condense and obtain ns-long coherence leading to $\gamma_{\text{pol}} \ll \Omega_{\text{M}}$. Following this, I will focus on the experimental phenomena related to this non-adiabatic interaction regime, such as tunable phonon sidebands in the emission. I will show that polariton-phonon interaction rate can be controlled by tuning the phonon population by transducers. Polaromechanical MCs are a powerful platform to study phonon lasing [6], strong polariton-phonon coupling [7], locking of polariton lasers by phonons [8], as well as applications in microwave-to-optical conversion.

- [1] P. V. Santos and A. Fainstein, *Polaromechanics: polaritonics meets optomechanics*, Opt. Mater. Express **13**, 1974 (2023).
- [2] J. Kasprzak et al., *Bose-Einstein condensation of exciton polaritons*, Nature **443**, 409 (2006).
- [3] A. Fainstein et al., *Strong optical-mechanical coupling in a vertical GaAs/AlAs microcavity for subterahertz phonons and near-infrared light*, Phys. Rev. Lett. **110**, (2013).
- [4] D. H. O. Machado et al., *Generation and propagation of super-high-frequency bulk acoustic wave in GaAs*, Phys. Rev. Appl. **12**, 1 (2019).
- [5] A. S. Kuznetsov et al., *Electrically Driven Microcavity Exciton-Polariton Optomechanics at 20 GHz*, Phys. Rev. X **11**, 021020 (2021).
- [6] D. L. Chafatinos et al., *Polariton-Driven Phonon Laser*, Nat. Commun. **11**, 1 (2020).
- [7] A. S. Kuznetsov et al., *Microcavity phononitons – a coherent optical-to-microwave interface*, Nat. Commun. **14**, 5470 (2023).
- [8] D. L. Chafatinos et al., *Asynchronous locking in metamaterials of fluids of light and sound*, Nat. Commun. **14**, 3485 (2023).